

SPECTROSCOPY OF THE HOST GALAXY OF THE GAMMA-RAY BURST 980703¹

S. G. DJORGOVSKI², S. R. KULKARNI², J. S. BLOOM²,
 R. GOODRICH³, D. A. FRAIL⁴, L. PIRO⁵, AND E. PALAZZI⁶

² Palomar Observatory, California Institute of Technology, Pasadena, CA 91125, USA

³ W. M. Keck Observatory, Kamuela, HI 96743, USA

⁴ National Radio Astronomy Observatory, Socorro, NM 87801, USA

⁵ Istituto di Astrofisica Spaziale, CNR, Via Fermi 21, 00044 Frascati, Italy

⁶ Istituto Tecnologie e Studio Radiazioni Extraterrestri, CNR, 40129 Bologna, Italy

Draft version February 1, 2008

ABSTRACT

We present spectroscopic observations of the host galaxy of the γ -ray burst (GRB) 980703. Several emission and absorption features are detected, making the redshift, $z = 0.966$, completely unambiguous. This is only the third known redshift for a GRB host. The implied isotropic γ -ray energy release from the burst is in excess of 10^{53} erg, for a reasonable choice of cosmological parameters. The spectroscopic properties of the host galaxy are typical for a star formation powered object. Using the observed value of the Balmer decrement, we derived the extinction in the galaxy's restframe, $A_V \approx 0.3 \pm 0.3$ mag. Using three different star formation rate indicators, we estimate $SFR \approx 10 M_\odot \text{ yr}^{-1}$, or higher, depending on the extinction, with a lower limit of $SFR > 7 M_\odot \text{ yr}^{-1}$. This is the highest value of the star formation rate measured for a GRB galaxy so far, and it gives some support to the idea that GRBs are closely related to massive star formation.

Subject headings: cosmology: miscellaneous — cosmology: observations — gamma rays: bursts

1. INTRODUCTION

Studies of the cosmic γ -ray bursts (GRBs; Klebesadel, Strong, & Olson 1973) are currently one of the most active fields of research in astronomy. Much of the recent progress in this area was enabled by the discovery of long-lived x-ray afterglows of GRBs (e.g., Costa et al. 1997), made possible by the BeppoSAX satellite (Boella et al. 1997). This, in turn, enabled the discovery of optical (e.g., van Paradijs et al. 1997, Bond 1997, Djorgovski et al. 1997, Halpern et al. 1998, Jaunsen et al. 1998, Hjorth et al. 1998, etc.), and radio (e.g., Frail et al. 1998, Taylor et al. 1998) afterglows and their subsequent studies, including the determination of their distances and various physical parameters.

The key measurement is the determination of redshifts. This has demonstrated convincingly the cosmological nature of GRBs (cf. Paczyński 1995), and determined the energetics scale of this fascinating phenomenon. To date, redshifts have been obtained for only two GRB host galaxies: $z = 0.835$ for GRB 970508 (Metzger et al. 1997, Bloom et al. 1998a), and $z = 3.428$ for GRB 971214 (Kulkarni et al. 1998). The two cases have implied energy releases which differ by more than an order of magnitude (Djorgovski et al. 1997, Ramaprakash et al. 1998, Kulkarni et al. 1998). Obviously, there is a pressing need to increase the size of this sample. Moreover, the physical properties of GRB host galaxies, especially the star formation rates, provide essential clues about the possible physical origins

of the GRB phenomenon.

Following the discovery of GRB 980703 by the RXTE satellite (Levine et al. 1998), a variable x-ray source was localised in its error-box by the BeppoSAX satellite (Galama et al. 1998a, Galama et al. 1998b). At the same time, radio observations from the VLA and optical observations from the Keck and Palomar lead to the discovery of a radio and optical transient (OT) within the error-circle of the fading x-ray source (Frail et al. 1998a, Frail et al. 1998b, Bloom et al. 1998b). Bloom et al. 1998c report in detail on these observations. This was subsequently confirmed by Zapatero-Osorio et al. 1998 and Castro-Tirado et al. 1998. In this Letter we report on the spectroscopy of the host galaxy of GRB 980703. Our preliminary results have been reported in Djorgovski et al. 1998.

2. OBSERVATIONS AND DATA REDUCTIONS

Spectra of the host galaxy were obtained using the Low Resolution Imaging Spectrograph (LRIS; Oke et al. 1995) mounted at the Cassegrain focus of the Keck II 10-m telescope, on UT 1998 July 07.6 and 19.6, in photometric conditions, and with sub-arcsecond seeing. There was some moonlight on July 19. On July 7 we used a 300 lines mm^{-1} grating giving a typical resolution of $FWHM \approx 12 \text{ \AA}$, and a usable wavelength range from approximately 3800 to 8800 $\text{\AA}$. The total exposure was 1 hour. Exposures of the spectroscopic standard BD +17°4708 (Oke & Gunn 1983) were used for flux calibration. On July 19 we used an 831 lines mm^{-1} grating giving a typical resolution of

¹ Based in part on observations obtained at the W. M. Keck Observatory, which is operated by the California Association for Research in Astronomy, a scientific partnership among the California Institute of Technology, the University of California, and the National Aeronautics and Space Administration.

$FWHM \approx 4 \text{ \AA}$, and a usable wavelength range from approximately 7900 to 9800 $\text{\AA}$. The total exposure was 0.5 hours. Exposures of the spectroscopic standard BD +28°4211 (Massey et al. 1988) were used for flux calibration.

Exposures of arc lamps obtained right after the target observations were used for the wavelength calibration, with a resulting r.m.s. uncertainty of about 0.3 $\text{\AA}$ for the July 7 data, and 0.1 $\text{\AA}$ for the July 19 data, and with the possible systematic errors of the same order, due to the instrument flexure. All observations were obtained with the slit position angle close to the parallactic angle, so that the atmospheric refraction effects were minimized. The estimated flux zero-point calibration is less than 10%. We tested the flux calibration consistency of the two spectra by comparing the line fluxes of two emission lines detected in both observations (identified as $H\delta$ and $H\gamma$; see below); these fluxes should remain constant, while the continuum intensity was expected to change, due to the fading of the OT. The line fluxes agree to better than 10%.

The resulting spectra are shown in Figs. 1 and 2. Several strong emission lines are observed, as listed in Table 1. The estimated errors of the observed line fluxes are less than 5% internal (due to the random noise and the uncertainties in the continuum level determination), plus about 10% external, due to the uncertainties in the flux calibration, including aperture corrections and other possible systematics. In addition, several absorption features are seen in the blue part of the spectrum, as shown in Fig. 3 and listed in Table 2. The mean emission-line redshift is $z_{em} = 0.9662 \pm 0.0002$, and the mean absorption redshift is $z_{abs} = 0.9656 \pm 0.0006$. The corresponding restframe velocity difference is $\Delta V = (90 \pm 110) \text{ km s}^{-1}$, which is consistent both with zero and with velocity fields expected in normal galaxies.

The observed continuum flux contains light from both the host galaxy and the OT. Our spectroscopic measurements give magnitudes $B \approx 23.3$, $V \approx 22.65$, $R \approx 21.9$, and $I \approx 21.3$ mag for the July 7.6 data, and $I \approx 22.0$ mag for the July 19.6 data, uncertain by about 0.2 mag. These are in a good agreement with the magnitudes estimated for these epochs from our direct imaging (Bloom et al. 1998c). We note that the OT accounted for about one-third of the observed flux in the R and I bands on July 7, and the observed equivalent widths of emission lines should be corrected accordingly, i.e., relative to the galaxy's light they should be increased by about 50%. The contribution of the OT on July 19 was negligible.

We estimate the foreground Galactic reddening in this direction from the maps by Schlegel, Finkbeiner & Davis 1998 to be $E_{B-V} = 0.061$ mag. Using the Galactic extinction curve from Cardelli, Clayton & Mathis 1989, we obtain extinction values of 0.251, 0.188, 0.141, 0.090, 0.053, 0.036, and 0.021 mag in the $BVR IJHK$ bands, respectively. Table 1 also lists the extinction-corrected emission line fluxes.

3. DISCUSSION

This is only the third known redshift for a host galaxy of a GRB. Assuming a standard Friedman model cosmology with $H_0 = 65 \text{ km s}^{-1} \text{ Mpc}^{-1}$ and $\Omega_0 = 0.2$, for $z = 0.9662$ we derive a luminosity distance $d_L = 1.921 \times 10^{28} \text{ cm}$.

The observed γ -ray fluence of this burst was $(4.59 \pm$

$0.42) \times 10^{-5} \text{ erg cm}^{-2}$ (Kippen et al. 1998; cf. also Amati et al. 1998). The corresponding isotropic energy release for our assumed cosmology is $(1.1 \pm 0.1) \times 10^{53} \text{ erg}$, making it nearly as luminous as GRB 971214 (Kulkarni et al. 1998, Ramaprakash et al. 1998), and thus showing that GRB 971214 was not peculiar in this regard.

The low-ionization emission-line spectrum of the host galaxy of GRB 980703 is typical of an actively star-forming galaxy. There is no sign of an active nucleus, given the absence of any high-ionization lines, and the relative weakness of the $[\text{O III}] 4959$ line compared to $H\beta$.

The observed equivalent width of the $[\text{O II}] 3727$ line on July 7 was $(91 \pm 5) \text{ \AA}$, or about 46 $\text{\AA}$ in the galaxy's rest-frame. Correcting for the estimated continuum contribution from the OT would increase these numbers by about 50%, placing this object at the high end of the distribution for the typical field galaxies at comparable magnitudes and redshifts (Hogg et al. 1998).

Given that we observe three lines of the Balmer series, we can make an estimate of the effective extinction in the galaxy's star forming regions. We observe line ratios $H\gamma/H\beta = 0.46 \pm 0.01$ and $H\delta/H\beta = 0.23 \pm 0.01$ (note that all systematic errors in flux calibration cancel for these ratios). We assume that the fractional correction due to the fill-in of the stellar absorption lines in the spectrum is the same for all three Balmer lines. The corresponding ranges for the standard Case B recombination (Osterbrock 1989) for a broad range of temperatures and electron densities are $H\gamma/H\beta = 0.469 \pm 0.009$ and $H\delta/H\beta = 0.229 \pm 0.004$. The ratios of the observed and theoretical values are thus 0.98 ± 0.04 and 0.89 ± 0.06 , respectively, indicating that some extinction may be present at the source, which is not surprising. Using the Galactic extinction curve (Cardelli, Clayton & Mathis 1989) and the foreground screen approximation, the implied restframe extinction values are $A_V \approx 0.125 \pm 0.25$ mag from the $H\gamma/H\beta$ ratio, and $A_V \approx 0.47 \pm 0.28$ mag from the $H\delta/H\beta$ ratio, with the weighted average $A_V \approx 0.3 \pm 0.3$ mag (in the galaxy's restframe).

We can estimate the star formation rate in this object in three different ways:

The implied $[\text{O II}]$ line luminosity, corrected for the Galactic foreground, but not for any restframe extinction, is $L_{3727} = (1.41 \pm 0.15) \times 10^{42} \text{ erg s}^{-1}$. Assuming that the line is powered by star formation alone, the conversion from Kennicutt 1998, we estimate the star formation rate $SFR \approx (20 \pm 7) M_\odot \text{ yr}^{-1}$. Adopting our estimated restframe extinction of $A_V \approx 0.3$ mag would increase these numbers by 53%, i.e., to $SFR \approx 30 M_\odot \text{ yr}^{-1}$.

Our observed $H\beta$ line flux, corrected only for the Galactic foreground extinction, is $f_{H\beta} = (6.1 \pm 0.6) \times 10^{-17} \text{ erg cm}^{-2} \text{ s}^{-1}$. Using the theoretical, Case B recombination value of $H\alpha/H\beta = 2.85 \pm 0.1$ (Osterbrock 1989), we derive the predicted $H\alpha$ flux $f_{H\alpha} = (1.74 \pm 0.19) \times 10^{-16} \text{ erg cm}^{-2} \text{ s}^{-1}$. The implied line luminosity (again, not corrected for any restframe extinction) is $L_{6563} = (8.1 \pm 0.9) \times 10^{41} \text{ erg s}^{-1}$. Using the conversion from Kennicutt 1998, we estimate the star formation rate $SFR \approx (6.4 \pm 0.7) M_\odot \text{ yr}^{-1}$. Adopting restframe extinction of $A_V \approx 0.3$ mag would increase these numbers by 25%, i.e., to $SFR \approx 8 M_\odot \text{ yr}^{-1}$. Despite the extrapolation from $H\beta$ to $H\alpha$, this is probably a more reliable estimate than the star formation rate

derived from the density-sensitive [O II] line.

However, the observed Balmer line luminosities are likely to be underestimated, due to the underlying stellar absorption features at the same wavelengths. Depending on the mean stellar age and star formation history, the restframe equivalent width of the H β line *in absorption* may be in the range $\sim 5 - 10 \text{ \AA}$. Our observed equivalent width of the H β line in emission is $48 \text{ \AA}$, corresponding to $\sim 24.5 \text{ \AA}$ in the restframe. Thus the correction to the derived Balmer emission line luminosities may be as large as 20%, or higher. Correspondingly, our derived SFR should be adjusted upwards by such a factor.

Finally, an alternative estimate of the SFR can be obtained from the continuum luminosity at $\lambda_{rest} = 2800 \text{ \AA}$ (Madau et al. 1998). The observed continuum flux at the corresponding wavelength of $5505 \text{ \AA}$ in our July 7 spectrum is $\sim 3.1 \mu\text{Jy}$. Correcting for the Galactic foreground extinction, the value becomes $\sim 3.7 \mu\text{Jy}$. We estimate that the OT contribution at this time was about 30% of the total flux, and thus the flux from the galaxy itself to be about $2.6 \mu\text{Jy}$. The implied restframe power at this wavelength is then $P_{2800} \approx 6.13 \times 10^{28} \text{ erg s}^{-1} \text{ Hz}^{-1}$. Using the Madau et al. 1998 estimator, the corresponding star formation rate is $SFR \approx 7.8 M_{\odot} \text{ yr}^{-1}$, again without any restframe extinction correction, and in an excellent agreement with the estimate derived from the H β line. Adopting restframe $A_V \approx 0.3 \text{ mag}$ would increase these numbers by 70%, i.e., to $SFR \approx 13 M_{\odot} \text{ yr}^{-1}$.

We thus conclude that the conservative lower limit to the star formation rate is $SFR > 7 M_{\odot} \text{ yr}^{-1}$ (with no

restframe extinction correction at all), at more realistically $SFR \approx 10 M_{\odot} \text{ yr}^{-1}$ or even higher, depending on the actual extinction value.

We note that these estimates exclude any star formation which may be heavily obscured, and whose diagnostics would thus escape our detection. In this case one may expect to detect the dust heated by the newly formed stars (Hughes et al. 1998), or else the measure the non-thermal radiation, produced as a by-product of the ongoing star formation (Condon 1992). Unless our SFR is underestimated by an order of magnitude or more, we determined that both processes would be expected at a level of $\sim 10 \mu\text{Jy}$, and thus cannot be easily detected with current telescopes.

This star formation rate is higher than in either one of the two GRB host galaxies with previously determined redshifts (GRB 970508 and GRB 971214), but is still not exceptional. This finding gives some modest support to the idea that GRBs may be closely related to recent star formation (e.g., Totani 1997, Wijers et al. 1998).

We are grateful to the Directors of the W. M. Keck and Palomar observatories, Drs. F. Chaffee and W. Sargent, for the allocation of service observing time for the initial observations reported here, and to the staff of WMKO for their expert assistance with the observing runs, and to the entire BeppoSAX team for their efforts. This work was supported in part by the Bressler Foundation (SGD), and grants from the NSF and NASA (SRK).

REFERENCES

- Amati, L., Frontera, F., Costa, E. & Feroci, M. 1998, GCN Circ. No. 146
 Bloom, J. S., Djorgovski, S., Kulkarni, S. R., & Frail, D. A. 1998, ApJ, in press
 Bloom, J. S., Djorgovski, S. G., Kulkarni, S. R., & Frail, D. A. 1998b, GCN Circ. No. 136
 Bloom, J. S., et al. 1998c, ApJ, in press
 Boella G., et al. 1997, A&ASup, 122, 299
 Bond, H. E. 1997, IAU Circ. No. 6654
 Cardelli, J.A., Clayton, G.C. & Mathis, J.S. 1989, ApJ, 345, 245
 Castro-Tirado, A.J., et al. 1998, preprint
 ARAA, 30, 575
 Costa, E., et al. 1997, Nature, 387, 783
 Djorgovski, S., et al. 1997, Nature, 387, 876
 Djorgovski, S., Kulkarni, S. R., Goodrich, R., Frail, D. A., & Bloom, J. S. 1998 GCN Circ. No. 139
 Frail, D. A., Kulkarni, S. R., Nicastro, L., Feroci, M., & Taylor, G. 1997, Nature, 393, 35
 Frail, D. A., Halpern, J. P., Bloom, J. S., Kulkarni, S. R., & Djorgovski, S. G. 1998b, GCN Circ. No. 128
 Frail, D. A., Kulkarni, S. R., Bloom, J. S., & Djorgovski, S. G. 1998c, GCN Circ. No. 141
 Galama, T. et al. 1998a, GCN Circ. No. 127
 Galama, T. et al. 1998b, GCN Circ. No. 145
 Halpern, J. P., Thornstensen, J. R., Helfand, D. J., & Costa, E. 1998, Nature, 393, 41
 Hjorth, J., Andersen, M.I., Pedersen, H., Jaunsen, A.O., Costa, E., & Palazzi, E. 1998, GCN Circ. No. 109
 Hogg, D. W., Cohen, J. G., Blandford, R., & Pahre, M. A., ApJ, in press (astro-ph/9804129)
 Hughes, D. H., et al. 1998, Nature, 394, 241
 Jaunsen, A.O., Hjorth, J., Andersen, M.I., Kjærsmo, K., Pedersen, H., & Palazzi, E. 1998, GCN Circ. No. 78
 Kennicutt, R. C. 1998, ARAA, 36, in press (astro-ph/9807187)
 Kippen, R. M., et al. 1998, GCN Circ. No. 143
 Klebesadel, R. W., Strong, I. B., & Olson, R. A. 1973, ApJ, 182, L85
 Kulkarni, S. R., et al. 1998, Nature, 393, 35
 Levine, A., Morgan, E., & Munro, M. 1998, IAU Circ. No. 6966; and GCN Circ. No. 126
 Madau, P., Pozzetti, L., & Dickinson, M. 1998, ApJ, 498, 106
 Massey, P., Strobel, K., Barnes, J., & Anderson, E. 1988, ApJ, 328, 315
 Metzger, M., et al. 1997, Nature, 387, 878
 Oke, J. B., and Gunn, J. E. 1983, ApJ, 266, 713
 Oke, J. B., Cohen, J. L., Carr, M., Cromer, J., Dingizian, A., Harris, F. H., Labrecque, S., Lucinio, R., Schaal, W., Epps, H. & Miller, J. 1995, PASP, 107, 375
 Osterbrock, D. E. 1989, Astrophysics of Gaseous Nebulae and Active Galactic Nuclei, Mill Valley, CA: University Science Books
 Paczyński, B. 1995, PASP, 107, 1167
 Ramaprakash, A. N., et al. 1998, Nature, 393, 43
 Schlegel, D.J., Finkbeiner, D.P., & Davis, M. 1998, ApJ, 500, 525
 Taylor, G., Frail, D. A., Kulkarni, S. R., Shepherd, D. S., Feroci, M., & Frontera, F. 1998, ApJ, 502, L115
 Totani, T. 1997, ApJ, 486, L71
 van Paradijs, J., et al. 1997, Nature, 386, 686
 Wijers, R. A. M. J., Bloom, J. S., Natarajan, P., & Bagla, J. S. 1998, MNRAS, 294, 13
 Zapatero-Orsorio, M. R., et al. 1998, GCN Circ. No. 130

TABLE 1
OBSERVED EMISSION LINES

λ_{obs} ($\text{\AA}$)	Line ID	z_{em}	f_{obs} ^a	f_{corr} ^b
7327.9	[O II] 3727.4	0.9660	26.9	30.4
8065	H δ 4101.7	0.9663	1.3	1.45
8533.5	H γ 4340.5	0.9661	2.55	2.8
9558.1	H β 4861.3	0.9662	5.65	6.1
9751	[O III] 4958.9	0.9664	2.5	2.7

^a Observed fluxes in units of 10^{-17} erg cm $^{-2}$ s $^{-1}$.

^b Corrected for the Galactic extinction.

TABLE 2
OBSERVED ABSORPTION LINES

λ_{obs} ($\text{\AA}$)	λ_{vac} ($\text{\AA}$)	Line ID	z_{abs}
4605.9	4607.2	Fe II 2344.2	0.9653
4669.9	4671.2	Fe II 2374.5	0.9672
4683.6	4684.9	Fe II 2382.8	0.9662
5083.8	5085.2	Fe II 2586.7	0.9654
5109.9	5111.3	Fe II 2600.2	0.9657
5493.6	5495.2	Mg II 2796.4	0.9651
5508.1	5509.6	Mg II 2803.5	0.9653
5604.6	5606.2	Mg I 2852.1	0.9656

Fig. 1. The initial spectrum of the host galaxy of GRB 980703, obtained at the Keck telescope on 7 July 1998 UT, when the OT was still contributing about one third of the observed continuum. The spectrum was smoothed with a gaussian with a $\sigma = 5\text{\AA}$, roughly corresponding to the instrumental resolution. Prominent emission and absorption lines are labeled.

Fig. 2. The far-red spectrum of the host galaxy of GRB 980703, obtained at the Keck telescope on 19 July 1998 UT. The spectrum was smoothed with a gaussian with a $\sigma = 2\text{\AA}$, roughly corresponding to the instrumental resolution. Prominent emission lines are labeled. The contribution of the OT to the spectrum was negligible at this point.

Fig. 3. A zoom-in on the spectrum from July 19, with the prominent absorption features labeled. They are most likely caused by the ISM in the GRB host galaxy itself. The spectrum was smoothed with a gaussian with a $\sigma = 1 \text{ pixel} \approx 2.5\text{\AA}$.




FIG. 1.—

